

Overestimate of committed warming

ARISING FROM C. Snyder. Evolution of global temperature over the past two million years. *Nature* (2016); doi: 10.1038/nature19798

Gavin A. Schmidt¹, Jeff Severinghaus², Ayako Abe-Ouchi^{3,4}, Richard B. Alley⁵, Wallace Broecker⁶, Ed Brook⁷, David Etheridge⁸, Kenji Kawamura^{9,10,11}, Ralph F. Keeling², Margaret Leinen², Kate Marvel^{11,12}, Thomas F. Stocker¹³

¹ NASA Goddard Institute for Space Studies, New York, New York 10025, USA. (email: gavin.a.schmidt@nasa.gov)

² Scripps Institution of Oceanography, University of California, San Diego, La Jolla, California 92093, USA.

³ Atmosphere and Ocean Research Institute, University of Tokyo, Kashiwa, Japan.

⁴ Japan Agency for Marine-Earth Science and Technology, Yokohama 236-0001, Japan.

⁵ Department of Geosciences, Earth and Environmental Systems Institute, Pennsylvania State University, University Park, Pennsylvania 16802, USA.

⁶ Lamont-Doherty Earth Observatory, Columbia University, Palisades, New York, New York 10964, USA.

⁷ College of Earth, Ocean, and Atmospheric Sciences, Oregon State University, Corvallis, Oregon 97331, USA.

⁸ CSIRO Climate Science Centre, Aspendale, Victoria, Australia.

⁹ National Institute for Polar Research, Tachikawa, Tokyo 190-8518, Japan.

¹⁰ Department of Polar Science, SOKENDAI (The Graduate University for Advanced Studies), Tachikawa, Tokyo 190-8518, Japan.

¹¹ Institute of Biogeosciences, Japan Agency for Marine-Earth Science and Technology, Yokosuka 237-0061, Japan.

¹² Department of Applied Physics and Applied Mathematics, Columbia University, New York, New York 10025, USA.

¹³ Climate and Environmental Physics, Physics Institute, University of Bern, Sidlerstrasse 5, CH-3012 Bern, Switzerland.

Palaeoclimate variations are an essential component in constraining future projections of climate change as a function of increasing anthropogenic greenhouse gases¹. The Earth System Sensitivity (ESS) describes the multi-millennial response of Earth (in terms of global mean temperature) to a doubling of CO₂ concentrations. A recent study² used a correlation of inferred temperatures and radiative forcing from greenhouse gases over the past 800,000 years³ to estimate the ESS from present day CO₂ is about 9°C, and to imply a long-term commitment of 3-7°C even if greenhouse gas levels remain at present-day concentrations. However, we demonstrate that the methodology of ref. 2 does not reliably estimate the ESS in the presence of orbital forcing of ice age cycles and therefore conclude

that the inferred² present-day committed warming is considerably overestimated. There is a Reply to this Comment by Snyder, C. W. *Nature* 547, <http://dx.doi.org/10.1038/nature22804> (2017).

The previous analysis² was based on the assumption that greenhouse gases were solely responsible for long-term global-mean glacial– interglacial temperature changes. This is not correct^[4–8]. While it is clear that greenhouse gases played a significant role, quantification of that role is difficult because of simultaneous changes in many factors that also influence the energy balance of the Earth (e.g. the extent of the ice sheets, snow cover, vegetation, dust load, cloud cover)⁵. By contrast, it is widely accepted that orbital forcing is the ultimate trigger for glacial-interglacial temperature change^[4–8], enhanced by fast and slow feedbacks involving the ice albedo, clouds, the carbon cycle, vegetation, etc.¹, sometimes even resulting in hysteresis behaviour⁶. Hence, the strong correlation seen in the analysed datasets² is a conflation of the sensitivity of the climate to CO₂ and the response of the carbon cycle to temperature and ice sheet variations. The Charney climate sensitivity (including fast atmospheric feedbacks, but not including long term ice sheet and vegetation changes) can be constrained by these data by treating those long-term factors as forcings⁸. However, estimating the long-term sensitivity to greenhouse gas forcing alone requires constraints from periods that are not affected by the interaction of orbital forcing and ice sheets, or that include a model-based assessment of the response to other forcings^{1,9,10,11}.

To better illustrate the lack of connection between ESS and the scaled regression of temperature and greenhouse gas forcing, we use a simple three-component coupled model¹² for land ice, temperature and carbon that allows for orbitally-forced impacts on the ice sheets, as well as both short and long-term feedbacks to changes in carbon dioxide, and a two-way coupling between temperature and ice. Based approximately on differences between pre-industrial and the last glacial maximum⁴, we fix the response of CO₂ to temperature (20 ppm K⁻¹) and the radiative forcing related to ice (0.025 W m⁻² mSL⁻¹), and vary the non-Planck climate feedback and the ice sheet response to temperature in order to span a wide but plausible range of Charney and Earth System sensitivities. The system is driven by an external 80 kyr periodic signal applied directly

to the ice sheet component (Fig. 1a). We calculate the model ESS and the scaled regression over the glacial cycles (the linear slope in $\text{K W}^{-1}\text{m}^2$ multiplied by 3.7 W m^{-2}) and plot the ratio (Fig. 1b). If the latter were a good estimate of the response to CO_2 forcing alone, the ratio would be close to unity everywhere, but this is clearly not the case: biases are very large and pervasive. For the ranges of model parameters chosen the scaled regression is a significant overestimate of the actual model ESS value.

This shows that application of an aggregate regression from glacial periods, in which orbital forcing as well as greenhouse gases caused temperature variations, to the committed warming from current radiative forcing will therefore likely overestimate future warming. In addition, given the current estimate of the radiative imbalance¹², the future changes in vegetation and ice sheets that would be required as a response to current and committed short term warming to reach a 3-7°C warming would imply at least a doubling of the original forcing. Given the limited extent to which the current ice sheets can change areal coverage and the expected range of vegetation impacts, this seems implausible. Furthermore, the response to global forcing likely depends on the climate state^{1,13}. Any paleo-derived ESS must, if it is to be applied to the present day radiative imbalance, be defined in a way that estimates the impact of external radiative forcing only, and should be drawn from evidence from non-glacial base climate states.

In summary, we demonstrate that an ESS of $\sim 9^\circ\text{C}$ as defined² cannot be used to project future warming, and no evidence was presented to alter the most recent assessment of the present-day committed warming¹⁴.




Figure 1. a) Two examples of synthetic temperature and carbon dioxide cycles over 1 million years driven by an 80 kyr cycle in ice sheet forcing using a simple 3-component model for temperature, CO₂ and glacial ice (see Supplementary Information). b) The ratio of the scaled regression (the linear regression coefficient in $K W^{-1} m^2$ multiplied by $3.7 W m^{-2}$) to the model ESS over a range of plausible parameter space.

Supplementary Information

Model code and fuller documentation are available in the SI:

1. Simple-coupled-ice.pdf (PDF file)
2. iceage_cycles.R (R code, plain text)
3. Python notebook (ipynb file)

References

1. PALAEOSENS Project Members. Making sense of palaeoclimate sensitivity. *Nature* 491, 683–691 (2012).
2. Snyder, C. Evolution of global temperature over the past two million years. *Nature* 538, 226–228 (2016)
3. Bereiter, B. et al. Revision of the EPICA Dome C CO₂ record from 800 to 600 kyr before present. *Geophys. Res. Lett.* 42, 542–549 (2015).
4. Köhler, P. et al. What caused Earth's temperature variations during the last 800,000 years? Data-based evidence on radiative forcing and constraints on climate sensitivity. *Quat. Sci. Revs.* 29, 129–145 (2010).
5. Hays, J. D., J. Imbrie, and N. J. Shackleton, Variations in the Earth's Orbit: Pacemaker of the Ice Ages, *Science* 194, 1121–1132 (1976).

6. Raymo, M. E., L. E. Lisiecki, K. H. Nisancioglu, Plio-Pleistocene Ice Volume, Antarctic Climate, and the Global $\delta^{18}\text{O}$ Record, *Science* 313, 492-495 (2006).
7. Abe-Ouchi, A., et al. Insolation-driven 100,000-year glacial cycles and hysteresis of ice-sheet volume, *Nature* 500, 190-194 (2013).
8. Paillard, D., Quaternary glaciations: From observations to theories, *Quat. Sci. Rev.* 107, 11-24 (2015).
9. Hansen, J. et al. Target atmospheric CO_2 : Where should humanity aim? *Open Atmos. Sci. J.* 2, 217–231 (2008).
10. Lunt, D. J. et al. Earth system sensitivity inferred from Pliocene modelling and data. *Nature Geosci.* 3, 60–64 (2010).
11. Pagani, M et al. High Earth-system climate sensitivity determined from Pliocene carbon dioxide concentrations. *Nature Geosci.* 3, 27-30 (2009).
12. Johnson, G. C., et al. Ocean heat content [in “State of the Climate in 2015”]. *Bull. Amer. Meteor. Soc.* 97 (8), S66–S70.
13. von der Heydt, A. S., et al. Lessons on climate sensitivity from past climate changes. *Curr Clim Change Rep* (2016)
14. IPCC. Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA (2013)

Author Contributions:

G.A.S and J.S jointly conceived of this comment, G.A.S and K.D.M developed the simple model, and all authors contributed equally to the writing.